

Is the Continuous Glucose Monitor (CGM) Right for Me?

A CGM is a way to check your sugars all the time without needing to prick your finger. This Guide tells you about Continuous Glucose Monitors (CGMs) and helps you decide if a CGM is right for you.



What is glucose?
Do I have to test for my
blood sugar and now this
glucose thing too?

Glucose means
“sugar”. When you test
your blood sugar you are
testing your blood glucose.
CGM is a different way of
testing your glucose.



Acknowledgements

Anne Peters, M.D.

Martha Walker, RD, CDE

Valerie Ruelas, MSW

Sally McClain, MSW

Latino Health Access

Fran Milner, Illustrator

Lorena Sprager, Clear Language Specialist

Patient/Community Advisors: America Bracho, Sandra Betancourt, Edward F. Garcia, Rosy De Gloria Giraldo, Prado González, Ofelia Hernandez, Yadira Molina, Marina Perez, Jessica Porras, Carmen Rodriguez, Sara Serafin-Dokhan, Kristina Tyler, and Shawn K. Wahinehookae

Children's Hospital Los Angeles: Jennifer Baldwin, BSN, RN, CPN, CDE; Lelnaz Kashefipour, MPP; Shelly-Anne Baker, RN, CDE

CGM Companies: Abbott, Dexcom, Metronic, Senseonics

Funding provided by the Leona M. and Harry B. Helmsley Charitable Trust.

Dear Reader

We are a group of adults with type 1 diabetes. We gave advice on putting this guide together. We also gave advice on other guides you may want to read:

- Is the Insulin Pump Right for Me?
- How Do I Use an Insulin Pump?
- Is the Insulin Pen Right for Me?
- How Can I Manage My Type 1 Diabetes Better?
- Is the Continuous Glucose Monitor Right for Me?
- How Can a Continuous Glucose Monitor Help Me Manage My Diabetes Better?
- How Can a Continuous Glucose Monitor Help and Carb Counting Help Me Manage My Diabetes Better?

Some of us use a Continuous Glucose Monitor (CGM) and some of us do not. Some of us had thought about using CGM. Some had not thought about it until we helped with this guide.

We want to try and help people understand what we have learned from our teachers and from each other. That is why we want to share it with you.

CGMs have been around for many years. Lately they have become more accurate and easier to use. More people with diabetes use them than ever before. It is a very useful way to check your glucose, or sugar level. Many people like CGMs and some do not.

We want to help you figure out if you would like to use a CGM. And if you do, we offer ideas on how to use it.

We also hope that in the future there will be even more ways to help us manage diabetes and one day cure it. In the meantime, we invite you to be curious and explore the CGM.


Let's get started!

HERE IS WHAT VICTORIA HAS TO SAY ABOUT USING A CGM:

"I got my first CGM in 2019 and I love it. It helps me manage my diabetes better. I can see how different food and exercise affect my glucose. It is a tool that gives me information. I can make better decisions on how much insulin to give. I really like that I do not have to do so many finger sticks. I can also share my CGM data with my doctor."



I do not like checking my blood glucose all the time. Is there something else I can do?



I was tired of checking my blood glucose 8 times a day. I started using a CGM. Now I can see my glucose when I want and I can control my diabetes better.



I do not know about CGM... Does it hurt? Where do I wear it?

Introduction

I am curious to learn more about the CGM.

We are glad you are curious to learn more about the CGM. It could be right for you to manage your diabetes.

We want to help you learn about this awesome technology. Using the CGM to manage your diabetes can help improve your blood glucose, or sugar, control. It can give you more freedom.

Do you have more guides I could read? Yes. You may want to read our other guides:

- Is the Insulin Pump right for Me?
- How Do I Use an Insulin Pump?
- Is the Insulin Pen Right for Me?
- How Can I Manage My Type 1 Diabetes Better?
- Is the Continuous Glucose Monitor Right for Me?
- How Can a Continuous Glucose Monitor Help Me Manage My Diabetes Better?
- How Can a Continuous Glucose Monitor Help and Carb Counting Help Me Manage My Diabetes Better?

I am the one that needs to make choices about how I treat my Type 1 Diabetes.

That is right. This is about your health, your life, and your choice to become the most active member on your diabetes team.

In this guide, we talk about your “team.” That is because it often takes many people to help you take care of your diabetes.

The first part of your team is always you. Family members and friends you include in helping you with your diabetes are part of your team. Then you have your doctor. You may also see a diabetes educator, a nurse, nurse practitioner or physician assistant. You may see a dietitian and maybe a social worker or a psychologist. Your pharmacist and eye doctor are part of your team, and anyone else you want to include. These people are part of your diabetes care team. Each can help you take better care of your diabetes.

Thank you for inviting us to join your team!

Table of Contents

SECTION 1 – Continuous Glucose Monitors (CGMs): The Basics

- 1** Can You Tell Me More About Glucose Level and Testing For It?
 - 2** What Are the Ways to Test For Glucose Level? How Do They Work?
 - 3** How Many CGM Types Are There?
 - 3** Is There Anything That Would Make Me *Not* Be Able to Use a CGM?
 - 3** Can You Tell Me About the CGM With Three Pieces and a *Short Lasting* Sensor? and What Do the Pieces Do?
 - 5** Can You Tell Me About the CGM With Three Pieces and a *Long Lasting* Sensor? and What the Pieces Do?
 - 5** Can You Tell Me About the CGM With Two Pieces?
 - 5** How Does a CGM Check My Glucose?
 - 5** Will the CGM Glucose Number Be the Same As the Number I Get With a Finger Stick and Glucometer?
 - 6** If I Use the CGM, Do I Still Need to Check My Glucose With a Glucose Meter?
 - 6** Are All CGMs the Same?
 - 7** What Do CGMs Look Like?
 - 7** What Different Things Can the CGMs Do? That Is, What Features Do They Have?
 - 9** How Often Do I Wear the CGM?
 - 9** Where On My Body Do I Wear the CGM?
 - 9** What Does “Site Rotation” Mean?
 - 10** How Often Should I Rotate to a Different Site?
 - 10** How Do I Insert the CGM Sensor?
 - 14** What Are the Steps For Inserting a Long Term CGM?
 - 14** How Do I Set Up the Receiver Or Smartphone?
 - 15** What Do I Need to Do Each Time I Put in a New Sensor?
 - 16** What Glucose Information Can I See On My Receiver Or Smartphone?
-

Table of Contents

- 16** Tell Me More About the Lines On the Screen. What Do Those Mean?
 - 18** What Other Things Should I Think About to Help Me Decide If I Should Get a CGM?
 - 19** Can You Tell Me More About CGM Alarms?
 - 20** How Would I Handle the CGM in Water? Like When I Bathe Or Spend a Lot Time in the Pool Or in the Ocean?
 - 20** What Should I Know About Having Sex With My Partner If I Use a CGM?
 - 20** If I Need a Medical Test Like An X-Ray, a CT Scan Or An MRI, Would I Have to Remove the CGM?
 - 20** Will My Medical Insurance Cover the Cost of the CGM?
 - 21** What If the Amount I Have to Pay From My Pocket For the CGM System Is Too Much?
 - 21** Are There Other Costs That I Need to Know About the CGM?
 - 22** What Would a CGM System Cost If I Had to Pay For It Myself?
 - 22** What If I Buy the CGM System On My Own and Then Have Trouble Paying For the Sensors and Transmitters?
 - 22** Is There Any Way to Get a Discount If I Pay For the CGM Myself?
 - 22** What Is the Basic Process For Getting a Sensor?
 - 23** How Long Do CGM Systems Last?
 - 23** How Will I Be Able to Learn All the Details On How to Use a CGM?
 - 24** What If I Try the CGM and I Do Not Like It?
-

Table of Contents

SECTION 2 – How a CGM Can Help Manage Type 1 Diabetes Better

25 There Are Many Reasons That the CGM Can Help You Manage Your Diabetes Better!

26 Can The CGM Help Me Lower My A1C?

27 Is Using a CGM Worth the Trouble of Learning How to Use It?

28 How Else Can I Learn About the CGM?

SECTION 3 – Conclusion

29 Am I Ready For a CGM?

30 Should I Rush Into This?

30 What Is the Next Step?

30 The Control of My Diabetes Is in My Hands!

30 Either Way, I Am Committed to Taking Care of My Diabetes.

APPENDIX 1 – Resources

APPENDIX 2 – Glossary of Diabetes Terms

SECTION 1 — Continuous Glucose Monitors (CGMs): The Basics

Can you tell me more about glucose level and testing for it?

Glucose level means sugar level. In this guide, we call sugar “glucose”. People with diabetes have to know what their glucose levels are to help them decide how much insulin to take or how much food to eat.

When someone does not have diabetes, an organ in the body called the pancreas makes insulin for the body. When the person eats it makes more insulin. When the person is not eating, it makes less insulin. The pancreas responds right away to the level of glucose in the blood so it knows how much insulin to make.

When someone has Type 1 Diabetes, the pancreas doesn’t make much insulin. So, when someone has Type 1 Diabetes, they need to give insulin with shots or the pump, just like a normal pancreas would do. For a person to know how much insulin to give, they need to know what their glucose level is.

Glucose level means how much glucose is in the blood. In the United States, this level is measured in milligrams per deciliter, or mg/dl. In other countries, it is measured in millimoles, or mmol/l.

Glucose levels go up and down. The main reason for this has to do with what a person has eaten, or how much insulin is in their body.




Glucose can be tested from either the blood or the interstitial fluid. The CGM tests interstitial fluid, which is in tissue just below the skin. We will tell you more about that on page 5.

A normal range (for someone without diabetes) is about 70 to 100 mg/dl (3.9–5.6 mmol/l) before breakfast and below 140 mg/dl (7.8 mmol/l) after meals. A person with diabetes does not have a normal range because their pancreas does not work. Their body does not make insulin. They must give insulin for the food they eat. If someone has diabetes, they will give more insulin if their glucose levels are high and less if they are low. If a glucose level is too low, a person will eat or drink something to raise their glucose. If a glucose level is too high, a person will give insulin. You must test your glucose level to know how much insulin to give.

What are the ways to test for glucose level? How do they work?

Glucose meter:

It used to be that the only way to test glucose level was to prick the skin. That is, squeezing some blood out from the finger tip onto a test strip. Then seeing the level on the glucose meter.




The glucose meter gives information on what the glucose level is at that moment the blood is tested. It does not say whether the glucose is going up, going down or staying the same. The glucose meter does not have alarms to alert someone to a level that is too low or too high.

Continuous Glucose Monitor (CGM):

A CGM is a very different way of testing the glucose level. A person puts it on a different part of their body every 6 to 14 days. After 6 to 14 days, a person throws out the one they have been wearing and then puts a new one on their body. There is also one type of CGM that can stay in the body up to 6 months.

The CGM checks the glucose is checked every few minutes. A person can now see their glucose level on a receiver or their smartphone without pricking their finger. And the CGM tells the person if their glucose level is going up, going down or staying the same. A CGM is a much easier way to check glucose levels and it gives a lot more information!



How many CGM types are there?

Some CGMs have three pieces. Others have two pieces. Most CGMs have a short lasting sensor. One has a long lasting sensor. A sensor is the part of a CGM that is under the skin and measures glucose.

CGM systems change and grow over time. Newer models are coming out all the time. Many become easier to use and wear. Others add new features.



Is there anything that would make me not be able to use a CGM?

Yes. If you have a severe tape allergy, you most likely will not be able to use a CGM. This is because you need tape to keep parts of the CGM in place.

Can you tell me about the CGM with three pieces and a short lasting sensor? And what do the pieces do?

Three piece CGMs have a **sensor**, **transmitter** and **receiver**.

The Sensor goes on the belly or arm, in most cases. It depends on the type of CGM. The sensor comes pre-loaded into a device that injects it under the skin. The device that puts the sensor under the skin is called an inserter. Once under the skin, the inserter is thrown away. The sensor stays in place for 10 to 14 days.




The sensor has a thin wire about a half an inch long that goes under the skin. The wire is attached to and covered by a plastic part. The plastic part of the sensor has tape on the bottom of it so that it sticks to the skin. This sensor part of the CGM is flat and about the size of an adult thumb print. Most CGM, sensors need to be replaced every 6 to 14 days.

The sensor tests the glucose level every few minutes. It then changes the information into an electrical signal. This signal goes to the transmitter.

The Transmitter is a small device that fits over the sensor. You attach it to the sensor. When you need to change the sensor, you take off the transmitter. You will be able to reuse the transmitter by placing it on a new sensor. So, do not throw it away!

The transmitter gets the signal from the sensor. It then sends the information to a receiver or a smartphone.

The sensor and transmitter are often taped down together to hold them in place.



The Receiver is the part that gets the signal from the sensor and transmitter so you can see your glucose levels. The receiver that comes with the CGM is about the size of a small smartphone.

There are options for how to get your glucose information. You can choose to use the receiver that comes with the CGM, or you can use your own smartphone. Some people use both the CGM receiver and their smartphone.



Some CGMs can send information some pump receivers. If you use an insulin pump, you can look at the pump receiver to check glucose levels.

You can see many helpful things on the receiver or smartphone screen. The most important thing is that you are able to see your glucose level as a number. You can also see arrows that show if your glucose is going up, going down or staying steady. You can send your glucose levels to the smartphone of a family member or friend if you want to.

If you use a smartphone, you will have to download an app to see your glucose reading. We will talk more about what else you can see on the screen on page 7

The receiver needs to be close by to receive a transmission. Some devices have a range of 20 feet. Others must be much closer — about 6 feet. You also have to make sure there is not anything between you and the receiver that will block the signal. For instance, it cannot be blocked a wall.

Can you tell me about the CGM with three pieces and a long lasting sensor? And what the pieces do?


The main difference between systems with a *short lasting* sensor and a *long lasting* sensor is how you place the sensor on the body. The long lasting sensor requires a small surgery to put a sensor under the skin. The sensor stays in place for up to 90 days. There is a transmitter that you have to place on top of the skin. Then the signal is sent to a smartphone.

Can you tell me about the CGM with two pieces?

Some systems have only two pieces. This is because the sensor and transmitter are one single piece. So, you do not have to attach the transmitter to the sensor. When you throw out the sensor, you get rid of the transmitter at the same time. You replace the combo sensor and transmitter every 10 to 14 days, depending on the brand. The sensor and transmitter unit send glucose information to a receiver or smartphone. The receiver for a two piece CGM is called a reader. At the moment, the two piece CGM does not have a way to send information to a pump receiver.

How does a CGM check my glucose?

The sensor sits under the skin. It reads the glucose level in something called the “interstitial fluid”. Interstitial fluid surrounds all the cells in the body. Interstitial fluid can also be called glucose fluid.



Will the CGM glucose number be the same as the number I get with a finger stick and glucometer?

The CGM and finger stick readings will not always be the same. The glucose reading from the CGM or finger stick shows you a number just from that moment in time. Glucose moves from the blood vessel into the tissue just below your skin. It takes 5 to 10 minutes for the glucose to move from the blood vessel to the tissue just below the skin. So, the CGM reading is showing what your glucose was 5 to 10 minutes ago.

The difference between the CGM and the finger stick is greater when your glucose is changing quickly. Glucose levels will change quickly right after you eat or give yourself insulin.

If you choose to use a CGM, you can learn how to use the feature that shows if your glucose is going up or down and how fast the glucose is changing. A finger stick reading does not give you all this information.

If I use the CGM, do I still need to check my glucose with a glucose meter?

It depends. Each CGM brand has different rules about checking blood glucose. Some devices require you to do finger sticks once or twice a day to calibrate the CGM. Calibrate means to enter your blood glucose number to help the CGM be more accurate. Others do not require any routine finger sticks. If you have a device that requires a blood glucose check, it will ask you to do it!

But all CGM brands suggest that you carry a glucose meter with you so you can test your fingerstick glucose level if for some reason the CGM number seems off. Also, sometimes the CGM stops working without warning. So it is always good to have a back up meter for glucose testing.

Are all CGMs the same?

No. Different companies make different CGMs. Right now in the United States, 4 companies have CGMs that are approved by the Federal Drug Administration (FDA). Newer and better CGMs are likely to be FDA approved in the future.

Each company makes a device with common parts. But each CGM has different features. Some examples of features are:

- How long a sensor lasts before it has to be changed
- If you have to do a finger stick to calibrate the CGM
- If it is approved for children to wear
- If it has alarms to tell you if your glucose is too high or too low

Here is a list of the CGMs approved in the United States:

COMPANY	CGM NAME	TYPE	APPROVED AGE
Abbott	Freestyle Libre Freestyle Libre 2	2 piece	18 and up 4 years and up
Dexcom	Dexcom G-5 and G6	3 piece	2 years and up
Medtronic	Guardian Connect	3 piece	14 to 75 years old
Sensionics	Eversense	2 piece	18 and up

What do CGMs look like?

Here are some photos of CGM systems:

FREESTYLE LIBRE



DEXCOM G-6



GUARDIAN CONNECT



EVERSENSE



What different things can the CGMs do? That is, what features do they have?

New and improved CGMs are being developed all the time. So, what is offered today might be different next year. Here are some things to think about when choosing a CGM.

All CGMs have:

- A small sensor that goes under your skin
- A way to insert the sensor
- A transmitter that gets information from the sensor and sends it to a receiver
- A receiver option so you can see your glucose levels on a screen or display
- A way to download or send the information to a software program that can make a report of what your glucose has done over a period of time

CGMs have different features:


- **Calibration:** Some CGMs need to be calibrated. This means taking a blood glucose reading from a fingerstick. Then you enter your blood glucose number on the meter into the receiver or CGM app on a smartphone. This helps to check that the CGM system is working as it should. Each CGM system has its own rules about how often you need to do finger sticks.
- **Age to be able to use the CGM:** The Federal Drug Administration (FDA) decides how old a person needs to be to use a CGM. Right now, only two systems are approved for young children.
- **Sensor life:** This is how long a sensor should stay under your skin. Most sensors can be used for 6 to 14 days. There is a long lasting sensor that stays under the skin for about 90 days.
- **Sensor warm-up time:** The warm up time is how long it takes for you blood glucose reading to be correct. Some CGMs take longer than others.
- **Transmitters either have batteries or can be charged:** You may need a new transmitter every 3 months. The receiver or your smartphone can also be recharged and these types of systems last years.
- **Display options:** This is how you see your glucose readings. Some CGM systems have their own handheld device called a receiver or scanner. Some can send information to a computer, smartphone or smartwatch. There are two systems that can send information to an insulin pump.
- **Alerts:** If your blood glucose is too high or too low, most CGMs can let you know. Some vibrate or let you know with an alarm sound. You can also customize or set the alerts the way you want them. One system does not have alarms.
- **Live data sharing:** This is a feature where you can share your blood glucose information with someone. They can see your information in real time the same way you can. Some people like to share data with their diabetes care provider to help with diabetes care decisions. Parents like to see what is happening with their children when they are apart.
- **Software and data analysis:** The information from the sensor can be sent or uploaded to software programs. Software programs take the information, or data, from the sensor. The programs analyze the data or turn the data into reports. These reports can be shared with your diabetes care provider. You can also see your own progress too.
- **Costs:** If you have insurance, you will have to check with the insurance company about which CGM systems they cover. You will also have to ask if they cover *all* costs. If you are paying, there are a few costs to consider. There are separate costs for sensors, transmitters, and receivers. For more information on costs see page 21.

How often do I wear the CGM?

All the time. The CGM sensor and transmitter are attached to your body at all times, even when you are sleeping.

You live your life and do all your activities wearing your CGM. You keep it on all the time. This includes:

- When you exercise
- When you take a shower
- When you go swimming
- When you wear a fancy outfit
- When you have sex



Where on my body do I wear the CGM?

Each CGM system has information on where to place the sensor on the body. The approved sites are the best. But people do find other spots that work well for them. People wear them in many places. This includes on the arms, abdomen or lower back and upper buttocks. You need to check your CGM manual to be sure of what the best places are to wear your sensor.

Most common places people attach the sensor is to the abdomen or back of the arm. This is because it is easy to reach the fatty tissue. People attach it above or below the beltline and waistline. It is best to avoid areas where the skin is puffy from putting sensors in the same place a number of times. You will learn where on your own body a sensor is best.

What does “site rotation” mean?

Site rotation means changing the place you put the sensor each time. This is so you do not put it in the same place every time. This is done to keep the skin and part under the skin healthy. If sensor place is not changed, lumps or hard areas can start to form under the skin. A sensor then can't work as well. Check your CGM manual for the best places on your body to put the sensor.

How often should I rotate to a different site?

Your diabetes team will tell you how often to move the sensor from one place to another. Most people need to replace and move, or rotate, the *short term* sensor every 6 to 14 days. There is one *long term* sensor that lasts 90 days.

Sometimes the diabetes team may tell you to rotate the site more often. They mostly do this if there is any sign the sensor site is not working right. If the sensor falls off before the time to wear it is up, you will need to call the company who makes the sensor to ask for another one.

How do I insert the CGM sensor?


You will learn how to put in the CGM sensors by yourself. This is the case unless you are using the Eversense CGM. Each CGM is a bit different. So you will have to follow your CGM brand instructions.

In general, sensors come with an insertion device that allows you to easily place the sensor. You will have to prepare to insert sensor first.


1. Check the sensor to make sure it has not expired.
2. Wash your hands.
3. Follow the instructions on package as to how to peel any covering. There is often a paper cover over the sticky part of the sensor tape. The sticky part will go on the skin.

Here are the basic steps to insert a sensor with 3 piece CGMs:


1. Clean your skin.




2. Place the inserter on the skin where you want the sensor to go.
 - a. Make sure it is about 3 inches from pumps sites
 - b. Avoid boney areas like your ribs
 - c. Do not use the same site two times in a row




- 3.** Push the button. It will make a popping sound. That lets you know the sensor is being inserted.



- 4.** Remove the inserter. The sensor will be in place.



- 6.** Attach the transmitter. Again, each CGM is a bit different. But in most cases you will put it on top of the sensor, press down and snap it in place.




- 7.** Some people use what is called over-tape. This is tape that you place over the sensor and transmitter to keep them in place.
- 8.** Now you are ready to set up the receiver.

Here are the basic steps insert a sensor for 2 piece CGMs:

CGMs with two pieces have the sensor and transmitter together as one piece:


- 1.** Check the single sensor and transmitter to make sure it has not expired.
- 2.** Wash your hands
- 3.** Prepare the inserter
 - a.** Follow the instructions on package as to how to place the sensor into the inserter. Open the sensor and inserter package. The next steps usually involve:
 - i.** Peel the lid off the sensor cup




- ii.** Line up the sensor and inserter mark



- iii.** Press down and twisting until you hear a click




- iv.** Lift the inserter out of the sensor cup.



- v.** The sensor is ready to be put on.



- 4.** Clean your skin.




- 5.** Place the inserter on the skin where you want the sensor to go.



- 6.** Push the button. It will make a popping sound. That lets you know the single sensor and transmitter is being inserted.

- 7.** Remove the inserter. The sensor will be in place.




- 8.** Press down on the tape around the sensor to make sure it is well stuck to your skin.
- 9.** Some people use what is called over-tape. This is tape that will be placed over the single sensor and transmitter to keep it in place.
- 10.** Now you are ready to set up the receiver.


What are the steps for inserting a long term CGM?

The Eversense CGM device is the long term one. A trained doctor in a clinic inserts it. It is considered a minor surgery. Once it is inserted, you will set up your receiver. Your doctor will have to remove or replace the sensor after 90 days.

1. The doctor makes incision, or small cut.



2. The doctor creates space under the skin and inserts the sensor.



3. The doctor closes the incision.



How do I set up the receiver or smartphone?

Your CGM system will most likely come with its own receiver. But you may also choose to use your smartphone. Some CGM systems will work with an insulin pump receiver. Setting up the receiver means adding information into the receiver or smartphone. To do this, you start by turning on the receiver or smartphone. If you are using a smart phone, you will have to download the app for the CGM you have.

Each system will have a set of menus to follow. These menus can be set for English or Spanish. The first time you use the receiver or smartphone app you will follow the prompts to set:


- Date and time
- Target glucose range
- Alarms, in some systems
 - Set the Low Alert alarm to let you know if your glucose is too low
 - Set the High Alert alarm to let you know if your glucose is too high

You may have to reset your receiver if the batteries die. You may also have to reset it if you need to make changes to your target glucose range or alerts.

What do I need to do each time I put in a new sensor?

You will have to do a few things.

Connect or pair the transmitter with the receiver or app on a smartphone. This is done so the transmitter can send glucose data to your receiver or smartphone so you can see the data. Turn on the Bluetooth function on the receiver or smartphone.




Start or activate your sensor. To do this you may have to enter in a sensor serial number into the receiver or smartphone. Some systems require that you scan the sensor by holding a receiver or smartphone over the single sensor and transmitter.

Calibrate the system in some systems. This means you will need to get a finger stick glucose reading from a glucose meter. You will enter your glucose reading into the receiver or smartphone.

Wait for a warm-up period for the sensor to start sending accurate glucose levels to the transmitter and receiver. How long the warm up period depends on which system you have. It can last anywhere from 1 to 24 hours.

What glucose information can I see on my receiver or smartphone?


The most important information you will be able to see on your receiver screen is your glucose level. You will be able to see this as a number. You will be able to see it as a wave pattern over time with your glucose target range highlighted. And you will see an arrow letting you know if your glucose is going higher or lower.



Tell me more about the lines on the screen. What do those mean?

Every time the CGM reads your glucose level it creates a dot. The dots then connect to make a line. This line shows your glucose over time.

The Y axis is the vertical line of numbers on the left side of the graph. This is your glucose number. **The X** axis is the horizontal line on the graph that is at the bottom of the graph. This is the time of day.




Throughout the day many things will make the line will go up and down. For instance, when you eat, the line will start to go up. When you take insulin, the line will start to go down.


There is a colored area of the chart that shows the best place for your glucose wave to be. This is called the target range. Some people may have a wave right in the target range. Others may have a wave that always seems to be above the target range. And still others may have a wave the often goes below the target range.

A line that drops below the target range is showing low glucose, or hypoglycemia. A line that is above the target range is showing high glucose, or hyperglycemia. Some people may have a flatter looking line and others may have a very wavy line.


CGM RECEIVER OR SMARTPHONE WINDOW:




This is an example of a flatter line in the target range:




This is an example of a wavy line with low glucose, or hypoglycemia:



This is an example of a wavy line with high glucose, or hyperglycemia:



Here is an example with a wavy line with both high and low glucose:



What other things should I think about to help me decide if I should get a CGM?

There may be other things you want to think about before getting a CGM that we talk about below. Many people feel that CGM makes their diabetes easier to manage. Others do not.

The upside of using a CGM is...

- You do not have to do so many finger sticks.
- It is a tool that you and your diabetes care team can use to help lower your A1C and keep your glucose in your target range.
- You can use the CGM at different levels. You can use it for just glucose level readings all the way to adjusting your insulin base on glucose patterns.



I like the alarms.
I don't have to worry
so much when
I sleep.

Even if you just use the CGM for the glucose level readings, it gives you and your team much more information than just finger sticks. And that can help all of you manage your diabetes better.

- Having alarms can give you an alert or even wake you up from sleep when your glucose is heading into a dangerous range.

There can be a downside of using a CGM...

Sometimes
I feel like the
CGM is my mother.
Always after me to say
how bad I am doing
with my diabetes!



- Sometimes people get upset at seeing their glucose levels all the time. This can be frustrating if you do not know what to do when you see a high or low number.
- Glucose levels are often higher for longer after meals than people realize. This can lead to giving insulin too soon after the meal. This is called “stacking”.
- All of us like to feel successful. And it can sometimes feel like failure if glucose levels are too high or too low much of the time.
- The alarms and alerts can be going off often. This can wake you or your partner up from sleep. It can also be a bother when you are going out with other people or are at work.

Can you tell me more about CGM alarms?

It is important to work with someone on your diabetes team to help you understand what the CGM alarm is telling you and how to respond.

You can change the alarms on your CGM. Most CGM alarms can be changed from making a sound to just vibrating. You can also turn off many of the alarms.

You can work with your diabetes team to figure out which alarms you need at first. Then you can change them over time. You do not want to get upset with your CGM for the alarms sounding too often!

If your alarms
seem to be going off
all of the time, I can
help you pick the most
important alarms to
start with.



How would I handle the CGM in water? Like when I bathe or spend a lot time in the pool or in the ocean?

In general, CGM sensors and transmitters are waterproof in up to 8 feet of water for at least 30 minutes. So you can swim, shower or take a bath.

The receiver is not waterproof and will need to be put in a dry area. The CGM sensor will keep collecting information, but if your receiver is too far away, the information will not be transmitted. This means your receiver will not get glucose readings. There will be a gap and you may not be able to see what your glucose level was while you were in the water or too far away from your receiver.

When in water, your receiver will need to be closer to you than when you are out of water. This is because the signal does not transmit as well in water.

A little secret about a CGM is that you do not have to wear it all the time. You can take a break from wearing it to go to the beach, the pool or a fancy party. You can take a break for a day or a week or two. Then you can go back to testing your finger sticks. But, most people like the CGM so much they do not want to go without it.

What should I know about having sex with my partner if I use a CGM?

Having an open talk with your partner often fixes any concerns you may have. The sensor and transmitter are quite small and are rarely an issue. The most common issue brought up by CGM users is that the alarms may start going off during sex as glucose level goes down or up.


If I need a medical test like an X-ray, a CT Scan or an MRI, would I have to remove the CGM?

Sometimes yes and sometimes no. Ask your doctor or the person doing the test if you can leave your CGM on or not.

Will my medical insurance cover the cost of the CGM?

Not all insurance plans cover the cost of the CGM. Some do and some do not.

Ask someone from your diabetes team if your health plan will pay for your CGM. They might know because they have prescribed CGMs before.



Your diabetes team can help you find out if your insurance will cover a CGM.

If your health plan will pay for a CGM, you can call them to find out which CGM you could get. If the CGM you are hoping for is not covered, it is ok to ask for what you want. Sometimes a “no” for a certain kind of CGM can turn into a “yes” if you have good reasons for a specific type of CGM.

Another way to find out is to contact the CGM company. These companies want to sell you a CGM. So, if you give them your name and health plan information they will find out if the insurance you have will cover the CGM.

Many health plans, or insurance companies, pay most of the cost of a CGM. But not all the costs. If your plan does cover the CGM, then also find out how much money you would have to pay.

What if the amount I have to pay from my pocket for the CGM system is too much?

You do not have to buy a CGM if you feel you cannot afford it.

Sometimes health plans have something called a “deductible.” That means each year you must cover some part of the costs of your medical care. This often means that it is better to buy big items like a CGM transmitter and receiver later in the year, when you have already paid all that you must for the year.

Are there other costs that I need to know about the CGM?

Other costs depend on the system you choose.

- You may need more sensors.
- You may need more transmitters.
- You may want a receiver. Many CGM systems come with their own receiver.
- You may use a smartphone. This cost will not be covered by insurance.
- There are supplies you will have to purchase not covered by insurance such as:
 - Tape to help keep the sensor and transmitter in place
 - Carry cases to hold receivers

Most often, your doctor will need to fill out forms and send them to your insurance company. Your insurance company will use these forms to decide what CGM system is

covered and how much you need to pay. You will need to see your diabetes team on a regular basis. During your visits, they will complete the forms, so you can keep getting your sensors and transmitters.


What would a CGM system cost if I had to pay for it myself?

If you do not have health insurance, CGM systems cost between \$500 and \$1,500 to get started. Monthly supplies, including sensors, cost about \$200 to \$500. This does not include smartphone costs.

What if I buy the CGM system on my own and then have trouble paying for the sensors and transmitters?

You can go on and off the CGM. So, if there is a time when you cannot pay for sensors or transmitters, you can go back to checking your glucose level with finger sticks. Then you can return to the CGM when you can pay for sensors and transmitters.

I lost my insurance for a while and went back to finger sticks.



Is there any way to get a discount if I pay for the CGM myself?

Some CGM companies may offer a discount if you are paying for it yourself. You can talk to your diabetes team or ask the CGM company if they offer any discounts.

What is the basic process for getting a sensor?

1. Some sensors are a **pharmacy benefit**. This means your doctor has to order the CGM system at a pharmacy, most often one of the bigger pharmacy chains. This is the most easy way to do it. These sensors are often the Libre and, in some cases, Dexcom.
2. If a CGM is **not a pharmacy benefit** through your insurance, then it is covered under Durable Medical Equipment (DME). This works in a different way. Your diabetes care team fills out a form called a Certificate of Medical Necessity (CMN). This is sent to the company that makes the CGM device. The manufacturer may be able to fill the order. Or they will send it to a distributor who will send you the CGM. If this happens, they should send your diabetes care team another form to fill out. This will also include sending notes from your chart.

Making sure the forms are done is a pain. But you have to do what you have to do...



This way is much more of a hassle for everyone. It is very important that you know who your distributor is. This is because they are the people who will ship you your CGM. Also, they are the people you will need to bug to send out the forms so your doctor can help you.

Sometimes your health plan will give you a choice of distributors. It is often a good idea to call around to see who charges the least.

How long do CGM systems last?

How long a CGM system lasts depends on the one you use.

Sensors: Short term sensors last between 6 to 14 days. The long term sensor, the Eversense implant, lasts up to 90 days.

Transmitters can last 2 weeks, 3 months or one year. Some you can recharge.

Receivers will last a long time if you take care of them. They usually come with a 1 year warranty.

CGM technology is changing all the time. In the future, sensors and transmitters will last longer. Smartphone apps will have new features for CGMs. Sharing your data will get easier. Whole new systems that are even better than what is on the market now will be offered.


How will I be able to learn all the details on how to use a CGM?

All this can appear difficult. Know that when you start, someone from your diabetes team will help you learn how to use the CGM. You will follow the instructions in the book that comes with your CGM. After you have been shown how to use the CGM, you will do it yourself.

If you have questions, you can call the phone number in your CGM book for help. Or you can call someone on your diabetes care team if you need help. You can also read our guide, “How Can a Continuous Glucose Monitor Help Me Manage My Diabetes Better?” to learn more.

What if I try the CGM and I do not like it?

Many people feel that their quality of life is better by using the CGM. They feel like they are more in control. But if you try it and do not like it, you can always go back to finger sticks to check your blood glucose.



SECTION 2 — How a CGM Can Help Manage Type 1 Diabetes Better

There are many reasons that the CGM can help you manage your diabetes better!

If you decide to get a CGM, be sure to read our guides “How Can Continuous Glucose Monitor Help Me Manage My Diabetes Better?” and “How Can a Continuous Glucose Monitor and Carb Counting Help Me Manage My Diabetes Better?”

Here are some basics on how the CGM can help you manage your diabetes better:

- You can see your glucose level anytime without having to prick your finger.
- You can see how your glucose changes over a few hours or days.
- You can see patterns or trends in what your glucose does over a period of time.
- Seeing glucose levels in real time can help you make more informed decisions throughout the day about how to balance your food, physical activity, and medicines.
- You can share your CGM reports with your diabetes care team to help make diabetes treatment decisions.
- You can share your glucose levels with a family member or friend.
- You can get alarms and alerts, so you know if your glucose levels are going too high or too low.
- Some sensors communicate with insulin pumps and direct them to deliver insulin.



I sleep better knowing my glucose is at a normal range. If is lower than I want, I often have a snack to make sure it does not go low at night.

I like sharing my reports with my team because together we can make better decisions on how I can control my glucose.



A CGM allows you to see how your blood glucose is doing all of the time. You can see this information on a receiver or your smartphone in real time. So, you can make decisions that will help you adjust how much insulin you need. Your CGM can alert you if your glucose is going too high or too low. With this information, you can better avoid complications from severe glucose lows or highs. You can also download data on your computer, tablet, or smartphone to see patterns and trends in your glucose levels. And you can share these with your diabetes team. You can print out reports that show what your glucose was doing over a period of time. The reports can show a summary of several days and you can see specific days too. The shared information can help you and your diabetes care team make the best plan for managing your diabetes.

Your CGM can:

- Record dangerously low overnight glucose levels, which often go undetected
- Track high glucose levels between meals
- Show early morning spikes in glucose
- Assess how diet and exercise affect you
- Find out if your treatment plan works on a day-to-day basis
- Show you how often you are meeting your target glucose range
- Show how much your glucose goes up and down throughout the day — this is called your glucose variability

With this information, your diabetes team can:

- Decide how much insulin you should take
- Develop an exercise plan that is right for you
- Suggest the number of meals and snacks you need each day
- Correct the types and doses of medications you are taking

Can the CGM help me lower my A1C?

Yes. A CGM can help you reach your glucose and A1C targets. One way the CGM can help is that it gives you knowledge about what is happening to your glucose all day and all night.

With finger sticks, you only know what is happening right at the time you are checking. Many people with diabetes get tired of checking so often. It is also hard to check when you are busy or want to sleep.

In case you do not know, an A1C is a three-month average of your blood sugar levels. You should have it checked every 3 months. For most people the goal for an A1C is to be less than 7%.



By knowing what your glucose level is, you can adjust your insulin. Or you can eat something to keep your glucose level from being too high or too low. You can set alarms to alert you when your glucose is going too low or too high. This way you do not have to constantly wonder what is happening with your glucose levels. You can also share your CGM reports with your diabetes team and make smarter changes with your medications.

All of these things can help you maintain better glucose control, which can lower your A1C.

Is using a CGM worth the trouble of learning how to use it?

Yes! This is because you can use the CGM system at different levels. You can just use it at the simplest level or all the way up to the most complicated. A CGM can help you manage your diabetes better, even at the simplest level.

What are the levels at which I can use the CGM?

LEVEL 1:

You will learn how to put on your CGM. And you will learn to link it to your receiver, pump receiver, or smartphone. You will also learn to set up the basic features of the receiver or smartphone. This includes time, date, start sensor, and general glucose target range. You will be able to find the screen that shows your glucose level. On the screen, you will see what your glucose level is and if it is going up or down.

The benefit of Level 1 is that the CGM is reading a glucose level about every 5 minutes. Also, you do not need a finger stick each time you want to see your glucose level. This will help you make decisions. For instance, it will help you decide if you should eat or take some insulin.

This level alone can help improve how you manage your diabetes.

LEVEL 2:

You will learn to set up more receiver functions. For instance, you will learn to set a personal glucose range and set alarms for when your glucose goes too high or too low. This can help you manage your glucose levels because the CGM can give you an alert if you are out of your glucose target range.

When you go to the doctor, you can make sure you are wearing your CGM and have either the receiver or your smartphone with you. This way, your doctor can download your glucose data and review it with you. Having many days of glucose data can help you and your diabetes team improve your treatment plan, if needed.

For the first 6 months, I just used the CGM to check what my glucose level was doing.



LEVEL 3:

As you get more comfortable with the CGM, you can learn to understand your glucose trends. A glucose trend is what your glucose typically does on any given day. Since the CGM stores each glucose reading, you can see what your glucose level has done over time. For instance, you may notice that your weekend days are different from your weekdays.

Learning your trends can help you plan changes to what and when you eat. It can also help you as to how you give yourself insulin. Trends give you insight to how different foods, activities, stress and other factors affect your glucose.

You can also share your data remotely with other people you choose. This includes your diabetes care team. What this means is that other people can see what is happening with your glucose levels just like you can. This feature can be very helpful if you need extra support to help manage your diabetes.

It did not take long for me to be at a level 3. I like to see what my glucose does over time. I like that I can share my data with my diabetes team too.



LEVEL 4:

At this level you will be very comfortable with the CGM and understand the different menus and information. You can also notice trends better. You can see a more in-depth picture of patterns. This can inform you on how much insulin you need. It can also inform you about what changes in your behavior can improve your glucose levels.

If you have internet or Wi-Fi access, you can upload your data to software programs sponsored by the company that makes your CGM. There are also other sites that have software that can take your CGM data. This data gets changed into reports that you can print out. These reports can show you your daily glucose values, trends and patterns as well as other glucose statistics. All of this can help you and your diabetes care team make the best decisions to manage your diabetes.

Using a CGM at *any* level can help you manage your diabetes better. To learn more, you can also read our guide, “How Can a Continuous Glucose Monitor Help Me Manage My Diabetes Better?”

How else can I learn about the CGM?

A great way to learn about the CGM is from people who use it. Your diabetes team may be able to suggest people you can talk with. Or you can look on-line to see what other people write about the CGM.

SECTION 3 – Conclusion

Am I ready for a CGM?

We hope that the information we have shared has helped you start to learn what the CGM is and how it works. It is up to you to decide if you want to try using one. But, if you have a severe tape allergy, you probably will not be able to wear a CGM. This is because you need tape to keep parts of the system in place on your skin.

Here is a checklist to help you see if you are ready to try the CGM. You can check “Yes,” “No” or “Not Sure” for each line. Fill it out and talk with your diabetes team about it.

This checklist
can help you decide
if a CGM is right
for you.



AM I READY FOR THE CGM? CHECKLIST

Are You Ready To:	Yes	No	Not Sure
Work closely with your diabetes team? This means going in for visits as often as needed, talking on the phone or emailing each other.			
Complete and pass a diabetes and CGM training to show you are ready to use the CGM safely?			
Try to figure out and fix CGM problems?			
Go back to finger sticks and blood glucose monitoring if your CGM is not working?			
Change your CGM sensor when needed?			
Follow site rotation plans?			
Call your diabetes team or CGM company if you have a problem?			
Hear or feel the alarms when your glucose is too high or too low?			
Calibrate your sensor if and when needed? This means do a finger stick.			
Look at your monitor about 10 to 12 times a day and not just rely on alarms?			
Be willing to learn how to use the CGM data to help how you manage your glucose? Your healthcare team can help you to turn your data into action.			

Should I rush into this?

In most cases, there is no rush to start using CGM. So, you can spend time thinking about it.

If I start on the CGM, am I stuck with it?

Keep in mind that you can get a CGM system and try it. If you change your mind you can go back to finger sticks. It is not permanent. In fact, some people like to take CGM breaks for many reasons.

If you do get one, you should try it for a few months to see how you adjust to it. The first few weeks of wearing it will be new and it will take a while to learn how to use. Then after you have worn it for a while, you will begin to see if you like it or not.

What is the next step?

The next step can be to talk with your diabetes team to see if CGM is right for you. Ask to see a CGM to hold it and check it out. In most cases, insurance covers one or two types of CGM. So be sure to ask to see that type of CGM if you have insurance.

The control of my diabetes is in my hands!

That is right. No matter what you decide about the CGM, the control of your diabetes is in your hands. A CGM can help you control it with shots or a pump. When you control it, you can avoid many of the serious problems that can happen if your glucose stays too high or too low for too long.

Either way, I am committed to taking care of my diabetes.

Congrats for being committed to take good care of your diabetes. We know that it is not always easy to treat Type 1 Diabetes. We know you can do it well.

Keep up the good work!

APPENDIX 1 — Resources

In this appendix, we offer some resources that may help you. We have the information in this order:

- Organizations
- Carb Counting and Nutrition
- Insulin Pump Companies
- Insulin and Insulin Pens
- Diabetes Supplies and Medications

We have provided internet links. These are US based links so most information is in English. If a site is available in your language, we recommend using that link. To see the information in Spanish or other languages, try Google Translate. It does a pretty good job at changing the information into the language you wish. But, a computer program does this translation so it may not be accurate.

To use Google Translate go to <http://translate.google.com/manager/website/> and follow the step-by-step guide. This is free!

ORGANIZATIONS

AMERICAN ASSOCIATION OF DIABETES EDUCATORS (AADE)

(800) 338-3633

www.aadenet.org

This is a group of diabetes educators. This is their link for patient resources: <https://www.diabeteseducator.org/patient-resources>. Here they offer information for people with diabetes.

AMERICAN DIABETES ASSOCIATION (ADA)

(800) 342-2383

www.diabetes.org

Spanish link: http://www.diabetes.org/es/?loc=util-header_es

This is the biggest group of people with diabetes and diabetes professionals. It helps people with both Type 1 as well as Type 2 Diabetes. There are local chapters that you can contact for help or to volunteer. They also have an online store to buy books, gifts and other helpful items.

ACADEMY OF NUTRITION AND DIETETICS

(800) 877-1600

www.eatright.org

This is a site for information about food and nutrition.

CHILDREN WITH DIABETES

www.childrenwithdiabetes.com

This started as a group having mostly to do with children with diabetes. Now it includes young adults as well as parents of people with Type 1 Diabetes.

It is a good site to learn about many tools and resources for people with Type 1 Diabetes.

CLINICAL RESEARCH STUDIES WEBSITE

clinicaltrials.gov

This site lists all the clinical research studies that are in process in the United States. You can do a search using the key words “Type 1 Diabetes” if you want to find those research studies.

DIABETES MINE

www.healthline.com/diabetesmine

This is a blog about Type 1 Diabetes. It has been around for many years and is very helpful. A woman who has Type 1 Diabetes started it. It shares many people’s experiences and advice.

DIABETES SISTERS

www.diabetessisters.org

This is a group for women with diabetes, mostly Type 1. They share ideas and experiences.

DIABETIC DANICA

www.facebook.com/DiabeticDanica

Danica is a kind young woman with Type 1 Diabetes. She makes YouTube videos about having Type 1 Diabetes and how to use diabetes devices. These videos can be helpful.

DIABTRIBE

www.diatrIBE.org

DiatrIBE is a non-profit organization. It evaluates and comments on new approaches and treatments for diabetes, both Type 1 and Type 2.

GLU

www.myglu.org

GLU is the largest interactive on-line network for people with Type 1 Diabetes as well as their care givers and family members. The nonprofit Helmsley Charitable Trust funds it.

It offers excellent advice and information about Type 1 Diabetes. You can also connect with others who have the same questions and concerns about diabetes as you do.

JDRF

www.jdrf.org

This used to be the Juvenile Diabetes Research Foundation. It was started to help do research on Type 1 Diabetes. Now it helps people living with Type 1 Diabetes as well as funds research. There are local JDRF offices that may be helpful to you.

TRIAL NET

www.diabetestrialnet.org

This is a group of researchers who work on preventing and treating early Type 1 Diabetes. Contact your local Trial Net site for screening risk of new onset Type 1 diabetes for yourself or family members.

TUDIABETES

www.tudiabetes.org

Spanish link: <http://www.estudiabetes.org>

This is a large on-line group of people with both Type 1 and Type 2 Diabetes. They share concerns and ideas in Spanish about living with diabetes.

CARBOHYDRATE (CARB) COUNTING AND NUTRITION RESOURCES

CALORIE KING

www.calorieking.com

This offers information on foods, carbs, calories and more.

CARBS AND CALS

www.carbsandcals.com

This offers books and a \$5 App that gives pictures of foods and their carb count.

THE DIABETES CARBOHYDRATE AND FAT GRAM GUIDE

(The American Diabetes Association)

This guide has quick, easy meal planning using carbohydrate and fat gram counts. You can buy it on many shopping websites like Amazon, Barnes and Noble and the American Diabetes Association online store. www.store.diabetes.org

THE DOCTOR'S POCKET CALORIE, FAT & CARBOHYDRATE COUNTER

(949) 642-1993

Family Health Publications publish this. You can buy it on many shopping websites like Amazon and Barnes and Noble and the calorie king online store at www.calorieking.com.

FIGWEE

www.figwee.com

This is an iPhone App for \$2.99 that gives pictures of many different foods along with their carbohydrate count.

NUTRITION IN THE FAST LANE

(Franklin Publishing)

(800) 643-1993

www.fastfoodfacts.com

This book has nutrition information for 60 of the most common restaurants in the United States.

NUTRITION AND DIABETES

(International Diabetes Center)

(888) 637-2675

www.idcpublishing.com

This web site has books for sale in English and Spanish for \$3 on nutrition and diabetes.

INSULIN PUMP COMPANIES

These websites give you lots of information about their pumps. They also offer on-line lessons about how to use their pumps.

It can be very useful to look at these sites. You can learn about pumps. You can review how to use the pump you have as well.

ACCU-CHECK PUMPS: ROCHE DIAGNOSTICS

(800) 280-7801

www.accu-checkinsulinpumps.com

This site provides information on the Accu-check Spirit pump.

ANIMAS PUMPS: ANIMAS CORPORATION

(877) 937-7867

www.animas.com

These pumps include the Animas Ping and Animas Vibe.

OMNIPOD PUMPS: INSULET CORPORATION

(800) 591-3455

www.myomnipod.com

This site shares about the Omnipod system. It also gives you the option to try a demo Omnipod pump.

MINIMED PUMPS: MEDTRONICS, INC.

(800) 646-4633

www.medtronicdiabetes.com/home

This is the site for all the MiniMed Medtronic devices.

TSLIM PUMPS: TANDEM DIABETES CARE

(858) 366-6900

www.tandemdiabetes.com

This site describes the features of the TSlim pump.

INSULIN AND INSULIN PENS

SHORT ACTING INSULIN (REGULAR INSULIN) AND INTERMEDIATE ACTING INSULIN (NPH)

These are the oldest and lowest cost types of insulin. They are Regular insulin (short acting) and NPH insulin (intermediate acting insulin).

There are different names for these kinds of insulin including Novolin R, Humulin R, and others. Often these insulins come in vials. But sometimes they come in pens.

www.humulin.com/other-humulin-products.aspx

This offers information on Humulin Regular and NPH insulin as well as 70/30.

Novolin Regular and NPH do not have a website in the U.S. but you can buy them here.

www.diabetesselfmanagement.com/blog/relion-insulin-and-other-products-at-walmart

ReliOn Regular and NPH insulin come in vials. They are part of Walmart's low cost selection of diabetes supplies and products.

RAPID ACTING INSULIN

Apidra (Glulisine) made by Sanofi

www.apidra.com

These come in vials and pens.

Humalog (Lispro) made by Lilly Pharmaceuticals

www.humalog.com/index.aspx

These come in both disposable and refillable pens as well as vials.

Novolog (Aspart) made by Novo Nordisk

www.novolog.com

These come in both disposable and refillable pens as well as vials.

LONG ACTING INSULIN

Biosimilar Glargine

www.basaglar.com

This is a copy of the insulin known as glargine (U100 Lantus). It acts in a similar way and costs somewhat less. It is a long acting basal insulin.

U100 Lantus or Glargine insulin

www.lantus.com

This comes in vials and pens. It is a long acting basal insulin.

U300 Lantus or Glargine insulin

www.toujeo.com

This concentrated Lantus (glargine) insulin acts longer than U100 glargine. It only comes in a pen.

Levemir or Detemir insulin

www.levemir.com

Levemir comes in pens and vials. It is a long acting insulin but it is somewhat shorter acting than Lantus, Degludec or Toujeo.

Tresiba or Degludec insulin

www.tresiba.com

This is the very longest lasting basal insulin. It only comes in pens. It comes in two strengths: U100 and U200.

Glucagon Pens

www.lillyglucagon.com

This is the site for the Lilly brand of glucagon.

www.cornerstones4care.com/tracking/what-to-know/glucagen.html

This is the website for the Glucagon Kit which is the Novo-Nordisk brand of glucagon.

DIABETES SUPPLIES AND MEDICATIONS

CASES FOR INSULIN

www.frioinsulincoolingcase.com

These cases keep insulin cool and are easy to carry.

www.myabetic.com

These are carrying cases for insulin and supplies.

GLUCOSE TABLETS

www.dex4.com

These are one type of glucose tablet on the market. Many pharmacies have their own generic brands. You can look for a type of glucose tablet that you think tastes the best. But be warned, these do not taste like candy.

PEN NEEDLES

www.novonordisk.com/patients/diabetes-care/insulin-pens-and-needles.html

These are insulin pens and needles made by Novo-Nordisk.

PEN NEEDLES AND INSULIN SYRINGES

www.bd.com/diabetes

BD makes many diabetes products. They include syringes, pen needles and insulin infusion sets. BD offers very helpful educational information.

WEBSITE FOR COMPARING THE LOCAL COSTS OF MEDICATIONS

www.GoodRX.com

This is a good free App for finding the best prices for your medications. You enter the medication you are looking for and your location. Then it tells you the cost of it at your nearby pharmacies. It also gives you discount coupons.

APPENDIX 2 – Glossary of Diabetes Terms

In this glossary, we list and define key words that have to do with diabetes. You can use this to look up words you want to learn more about.

A1c

This is also:

- HbA1c
- Hemoglobin A1c
- Glycosylated hemoglobin

It is a blood test. The test can be a finger stick or blood taken from your vein. It tells you what your average blood sugar has been over the past three months. It does this by measuring the percentage of red blood cells in your body that have glucose stuck to them.

In most cases, normal A1c levels are 4% to 5.6%. The goal is to have your A1c as close to normal as possible, without having too many low blood sugar reactions. Your diabetes team will help you figure out what is the best target for you.

Be sure to do this test as often as your diabetes team orders it, about every 3 months.

Antibodies

These are proteins the body makes to protect itself from outside threats. These threats can include bacteria or viruses.

People get type 1 diabetes when their antibodies destroy the body's own beta cells that make insulin.

Aspart

This is the generic name of one kind of rapid-acting insulin. The branded (trade) name for aspart is Novolog. See rapid-acting insulin for more information.

Apidra

This is a branded (trade) drug name of one kind of rapid-acting insulin. The generic name for Apidra is glulisine. See rapid-acting insulin for more information.

Autoimmune disease

This is a disease caused by a problem in the body's immune (infection fighting) system that causes an attack on the body itself, rather than an infection. Type 1 diabetes is this kind of disease.

Basaglar

This is a brand drug name of one kind of basal insulin. The generic name for Basaglar is glargine. This long-acting basal insulin drug comes in one strength written as U100. See basal insulin and long-acting insulin for more information.

Basal insulin

You give this insulin with a shot once or twice a day. In most cases this insulin is only for patients on multiple daily shots. Basal insulin comes in different strengths shown as U100, U200 and U300. There are two types of basal insulin, long-acting insulin and intermediate acting insulin. See long-acting insulin and intermediate acting insulin for more information.

Basal insulins are:

Generic name	Brand name
NPH U100	Humulin (N) or Novolin (N) or ReliOn (N)
Degludec U100	Tresiba U100
Degludec U200	Tresiba U200
Detemir U100	Levemir
Glargine U100	Lantus or Basaglar
Glargine U300	Toujeo

Basal rate

Your body needs insulin on an ongoing basis even when you are not eating. The basal rate is the amount of insulin you need to give by shots or with an insulin pump. When the basal rate or basal insulin dose is set just right, the blood sugar does not go high or low when you are not eating.

For those using a pump, basal rates are in units per hour. You may see units per hour written as units/hour or u/hr. Typical rates are between 0.4 u/hr. and 1.6 u/hr. If you are using shots, you give yourself basal insulin doses in daily units, such as 15 units or 20 units. Your diabetes team will tell you what your basal doses should be.

Beta cells or β -cells

Beta cells or β -cells are cells that make insulin.

These cells are in the part of the pancreas called the Islets of Langerhans. See Cells for more information.

Blood glucose (BG) or Blood sugar

Blood glucose is also blood sugar.

This is the main sugar that is in the blood. This sugar is the body's main source of energy.

Bloodstream

The blood flowing through the circulatory system in the living body.

Blood sugar level

This means how much sugar is in the blood.

Blood sugar levels are measured in the U.S. in milligrams per deciliter, or mg/dl. In other countries, in milimoles, or mmol/l.

A normal range (for someone without diabetes) is about 70 to 100 mg/dl (3.9 to 5.6 mmol/L) before breakfast and below 140 mg/dl (7.8 mmol/L) after meals.

Blood sugar meter

This is a small, portable machine. People with diabetes use it to check their blood sugar levels.

After pricking the skin with a lancet, you place a drop of blood on a test strip. The test strip is placed in the machine. Then the meter, or monitor, shows the blood sugar level as a number on the digital display.

Blood sugar monitoring

This means checking your blood sugar level on a regular basis to manage diabetes.

You need a blood sugar meter or blood sugar test strips that change color when a drop of blood touches them. This is so you can check your blood sugar often.

Bolus

This is a burst of short or rapid acting insulin. It acts over a short period.

Most often, a bolus is to offset the blood sugar rise that happens after eating or drinking carbohydrates. It is also a correction dose to bring down a high blood sugar level back to normal.

The insulins for this are:

Generic name	Brand name
Insulin Regular	Humulin (R) or Novolin (R) or ReliOn (R)
Lispro	Humalog
Aspart	NovoLog
Glulisine	Apidra

Cannula

This is a small and flexible tiny piece of tubing. It stays under the skin once you remove the needle from the infusion set of an insulin pump.

Carbohydrate or Carb

Carbohydrates are also called carbs. Carbohydrates are one of the three main parts in foods:

1. Carbs
2. Fats
3. Proteins

They are the most important part of foods to control sugar. Carbohydrates are mainly sugars and starches. They have four calories per gram.

Carb bolus

This is a spurt of insulin that gets sent out quickly in the body to match carbs you are about to eat in a meal or snack. Most people use between 1 unit of rapid acting insulin for each 5 grams of carbs up to 1 unit for each 25 grams of carbs.

Carb counting

This means counting the grams of carbs in any food you eat or liquid you drink. This is a useful way to find out the amount of insulin you need to keep a normal blood sugar.

Carb factor or Carb Ratio or Insulin-to-carb ratio

This is the number of grams of carbs that one unit of insulin covers for a person. This varies from person to person. Your diabetes team will tell you your ratio.

Catheter

This is also pump tubing. Insulin goes through this plastic tube from the pump to the insertion set of a pump.

Cells

Cells are the smallest units of life. They are basic building blocks for all known life forms. Cells make up the parts of your body, like your skin, bones, heart, liver, or lungs. A person has over 10 trillion cells in their body.

Certified diabetes educator (CDE)

This is a health care professional with expertise in diabetes education. Trained and certified.

Continuous subcutaneous insulin infusion (CSII) or Insulin pump

CSII is the formal name for an insulin pump. See Insulin pump for more information.

Coma

This is a sleep-like state where a person is not conscious. Very high or very low blood sugar in people with diabetes can cause a coma.

Continuous glucose monitor (CGM)

A system consisting of a sensor, transmitter and receiver which determines subcutaneous or under the skin glucose levels every 1 to 5 minutes.

Correction bolus

A spurt of short or rapid acting insulin sent out quickly in the body. It is to bring a high blood sugar level back within a person's target range before a meal, after a meal, or at bedtime.

Correction factor or Insulin sensitivity factor

This is the fall in blood sugar level that one unit of insulin will produce. It is set by your diabetes team. It is often in the range of 25 to 75 but can be more or less depending on what your body needs.

A correction factor of 50 is used as a starting point. This means that 1 unit of insulin will lower your blood sugar by 50 mg/dl (2.8 mmol/L). For instance, if your correction factor is 50 and your blood sugar is 200 mg/dl (11.1 mmol/L), you expect that giving 1 unit of insulin will lower your sugar by 50 points. Which means that after 1 unit of insulin, the blood sugar will fall from 200 mg/dl (11.1 mmol/L) to 150 mg/dl (8.3 mmol/L).

Dehydration

This is when a person does not have enough water in their body. This can come from drinking too little fluid. It can also come from losing too much body fluid when a person pees or urinates often, sweats, has diarrhea or vomiting.

Delayed-onset hypoglycemia

A drop in blood sugar levels that can happen many hours after intense exercise.

Diabetes team

A group of people who help you take care of your diabetes. You are the most important member of your team. The other people on your team can be:

- Doctor
- Nurse or nurse practitioner or physician assistant
- Diabetes educator
- Dietitian or diabetes educator
- Social worker
- Psychologist
- Eye doctor

These people are part of your diabetes team. Each one of them can help you take better care of your diabetes.

Diabetic coma

This is when a person with diabetes is not conscious and is in a sleep-like state. Very high or very low blood sugar in people with diabetes can cause this.

Diabetic ketoacidosis (DKA) or Ketoacidosis

This is a very serious condition where the body does not have the insulin it needs. This results in dehydration and the buildup of acids in the blood. This needs to be treated in the hospital. It is life-threatening.

Dietitian

A health care professional who tells people about meal planning, carb counting, weight control and diabetes management. A registered dietitian (RD) has more training. Dietitians can also be diabetes educators.

Degludec

This is a generic drug name of one kind of basal insulin. The brand name for degludec is Tresiba. This long-acting basal insulin drug comes in two strengths written as either U100 or U200. See basal insulin and long-acting insulin for more information.

Detemir

This is a generic drug name of one kind of basal insulin. The brand name for detemir is Levemir. This long-acting basal insulin drug comes in one strength written as U100. See basal insulin and long-acting insulin for more information.

Endocrinologist

A doctor with the title MD or DO trained to treat diseases related to glandular problems. This includes diabetes.

Exchange lists

These lists are one of the ways for people with diabetes can plan meals. The lists have different types of food and show the amount carbs, proteins and fats in a serving size. Knowing this information helps you know how much insulin you will need if you eat that food.

Extended bolus

The insulin pump sends out a bolus over a fixed period set by the patient. For example, the pump could be set to give the bolus dose over 2 or 3 hours instead of right away. In most cases, the pump gives the bolus right away. This is a way to give insulin over a longer period, which is good for foods that the body absorbs more slowly, such as foods with a lot of fat in them.

Fasting

This means not eating food or drinking any fluids except water.

Fasting plasma glucose (FPG) test

A lab test that people take after fasting for 8 to 10 hours. In most cases, people fast overnight and take the FPG test in the morning.

An FPG level of less than 100 mg/dl (5.6 mmol/L) is normal. A level of 100 to 125 mg/dl (5.6 to 6.9 mmol/L) means prediabetes. A level of 126 mg/dl (7.0 mmol/L) or more means a person likely has diabetes. When a level is over 126 mg/dl (7.0 mmol/L), there will be more tests to confirm if the person has diabetes.

Fats

Fats are one of the three main parts of foods along with carbohydrates and protein. Fats occur alone as liquids or solids. This includes oils and margarines. They also can be a part of other foods.

Fats come from animals, veggies, nuts or seeds. Fats have 9 calories per gram.

Fiber

A kind of carb that passes through the digestive system intact. It does not raise blood sugar levels. It comes from plants.

Fiber adds bulk to your diet. It is very important for keeping your intestines healthy.

Food bolus

A dose of insulin that a person with diabetes takes before meals or snacks. This is to cover the expected rise in blood sugar from the food. Often, food boluses match the amount of carbohydrates in the food.

Glargine

This is a generic drug name of one kind of basal insulin. The brand name for glargine is Lantus or Basaglar or Toujeo. This long-acting basal insulin drug comes in two strengths written as either U100 or U300. See basal insulin and long-acting insulin for more information.

Glucagon—the hormone

This is a hormone. The alpha cells make it in the Islet of Langerhans in the pancreas. This hormone raises blood sugar levels. The opposite hormone to insulin that lowers blood sugar levels. In people without diabetes, the glucagon and insulin work together, to keep blood sugars normal. In people with diabetes, not enough glucagon is made to keep the blood sugars normal so they can fall too low.

Glucagon—the medication

Glucagon is given as a shot to help raise your blood sugar level. It is something that another person would give you if you were having a low blood sugar reaction and were not able to eat or drink sugar to bring it back up. The shot raises the blood sugar quickly. It does this by releasing sugar that is stored in the liver.

Glucagon emergency kit

A kit that has glucagon and a syringe. It is used to treat severe low blood sugar. Glucagon is a hormone that quickly increases blood sugar.

You need a prescription to get glucagon. It is a shot that someone else must give you. You should always have a glucagon kit at home, just in case. Be sure the one you have is not expired.

Glucose

A simple sugar that is in the blood. The body uses glucose for energy.

Glucose tablets

These are tablets that you chew and swallow. They are made of pure glucose. People take them to treat low blood sugar.

Glulisine

This is a generic drug name of one kind of rapid-acting insulin. The trade name for glulisine is Aprida. See rapid-acting insulin for more information.

Glycemic index (GI)

This is a method to classify foods, most of all carbs. The Index is based on how much the blood sugar level goes up after eating the certain food.

Glycogen

When you eat, carbohydrates they turn into a form of sugar called glycogen. This is a storage form of glucose in your liver and muscles. The glycogen is stored in your liver and muscles. When you have a low blood sugar, fast, or exercise, the glycogen turns into glucose and is release into the blood stream when you need it.

Gram

This is a small unit of weight in the metric system. People with diabetes use grams to weigh food.

Hormone

This is a chemical substance made by a gland or tissue. The blood carries it to other cells in the body. There, the hormone attaches to cells and causes them to do a certain job. For instance, when insulin attaches to a muscle cell it lets sugar go inside the cell. This is described as a “lock and key” effect. The hormone is the key and the cell is the lock. When the hormone insulin attaches to the cell, it opens the door and let’s sugar inside.

Insulin and glucagon are hormones.

Humulin [N]

This is a brand drug name of one kind of intermediate-acting insulin. The generic name for Humulin [N] is NPH. See intermediate-acting insulin for more information. It is a cloudy insulin.

Humulin [R]

This is a brand drug name of one kind of short-acting insulin. The generic name for Humulin [R] is Insulin Regular. See short-acting insulin for more information.

Humalog

This is a brand drug name of one kind of rapid-acting insulin. The generic name for Humalog is lispro. See rapid-acting insulin for more information.

Hyperglycemia or High blood sugar

This is when a person has a higher than normal level of sugar in the blood. In most cases, this means a blood sugar level of more than 180 mg/dl (10.0 mmol/L).

Hypoglycemia or Low blood sugar or Insulin reaction

This is when a person has a lower than normal sugar level in the blood. In most cases, this means a blood sugar level of less than 70 mg/dl (3.9 mmol/L).

Symptoms can vary. They can include feeling confused, nervous, shaky, drowsy or moody. They can also include sweating, headaches or numbness in the arms and hands.

If it is not treated, severe low blood sugar can cause loss of consciousness, convulsions, or even death.

Infusion set

This is part of an insulin pump. This set transfers insulin from the pump through an infusion line to below the skin. The set includes the tubing, tubing connector, insertion set, cannula and adhesive.

Infusion site or Insertion site

This is the area on the body where someone who uses an insulin pump inserts the cannula or needle.

Injection or Shot

This is when someone inserts liquid medication or nutrients into the body with a syringe. A person with diabetes injects insulin just under the skin, into what is subcutaneous tissue. Subcutaneous means below the skin.

Injection sites

These are places on the body where people most often inject insulin.

Injection site rotation and Infusion site rotation

The place you change on the body where you inject insulin or put the infusion sites. When you rotate, it prevents lipodystrophy. This means an abnormal build-up of fat under the skin.

Insertion set

The part of the infusion set that a person inserts through the skin. It may be a thin or a large metal needle. When the person removes the needle, it leaves a small Teflon catheter or cannula under the skin.

Insulin

This is a hormone made by beta cells in the Islet of Langerhans in the pancreas. The body sends out insulin when blood sugar levels go up, for instance after eating a meal. Its job is to lower blood sugar levels to normal.

Insulin lets sugar go into cells. Sugar gives your cells the energy to live. Without insulin, the sugar stays on the outside of the cells and goes up to very high levels in the blood. Without insulin, you would die because your cells would have no energy to live.

When your body cannot make its own insulin, there are different types for insulin drugs you can take. Your diabetes team will figure out the best insulin for you. The table below explains about the different types of insulin. You can also look up the types and names of insulin in this glossary for more information.

Generic Name (Brand Names)	Onset — Time for insulin to reach blood- stream	Peak — Period when insulin is most effective	Duration — How long the insulin works
RAPID-ACTING INSULIN			
Lispro (Humalog)	About 15 to 30 minutes	About 30 to 90 minutes	About 3 to 5 hours
Aspart (Novolog)	About 15 to 30 minutes	About 30 to 90 minutes	About 3 to 5 hours
Glulisine (Apidra)	About 15 to 30 minutes	About 30 to 90 minutes	About 3 to 5 hours
SHORT-ACTING INSULIN			
Insulin Regular [R] (Humulin [R], Novolin [R] or ReliOn [R])	About 30 minutes to 1 hour	About 2 to 5 hours	About 5 to 8 hours
INTERMEDIATE-ACTING INSULIN AND CALLED A BASAL INSULIN			
NPH [N] (Humulin [N], Novolin [N] or ReliOn [N])	About 1 to 2 hours	About 4 to 12 hours	About 18 to 24 hours
LONG-ACTING INSULIN AND CALLED A BASAL INSULIN			
U100 Glargine (Basaglar or Lantus)	About 1 to 1 and a half hours	Maybe slight peak at 12 hours in some people; no peak time in others	About 20 to 24 hours
U300 glargine (Toujeo)	About 1 to 1 and a half hours	No peak	About 28 to 36 hours
Detemir (Levemir)	About 1 to 2 hours	About 6 to 8 hours	Up to 24 hours
Degludec (Tresiba)	About 30 to 90 minutes	No peak time	About 42 hours
PRE-MIXED INSULIN			
	About 30 minutes	About 2 to 4 hours	About 14 to 24 hours
50% NPH/50% regular insulin Humulin 50/50	About 30 minutes	About 2 to 5 hours	About 8 to 24 hours
70% long acting/30% rapid acting insulin Novolog 70/30	About 10 to 20 minutes	About 1 to 4 hours	Up to 24 hours
75% long acting/25% rapid acting insulin Humalog mix 75/25	About 15 minutes	About 30 minutes to 2 and a half hours	About 16 to 20 hours

Insulin adjustments

A change in the amount of insulin a person with diabetes takes. Based on factors like meal planning, activity levels and blood sugar levels.

Insulin pen

A device that injects insulin. It looks like a pen for writing.

There are two kinds of insulin pens:

1. Prefilled pen with insulin that is disposable
2. Reusable pen that holds replaceable cartridges of insulin

To inject the insulin under the skin, you need to screw on a needle to the top of the pen.

Insulin pump

This is a small machine about the size of a small cellphone. It is computerized. You can program it to deliver a constant amount of basal insulin and give a bolus of insulin for a meal or high blood sugar. It takes the place of insulin shots.

A pump sends out fast-acting insulin through a plastic catheter, or tube. A Teflon infusion set or a small metal needle connects to the tube. You insert the set or small needle through the skin. The body gradually absorbs the insulin into the bloodstream.

Insulin Regular

This is a generic drug name of one kind of short-acting insulin. The brand name for insulin Regular is Humulin [R], Novolin [R], or ReliOn [R]. See short-acting insulin for more information.

Insulin sensitivity

This is a term to describe how the body reacts to insulin. Everyone reacts differently whether your body is making its own insulin or you must get insulin by shots or a pump. If a person is sensitive to insulin, it means that a smaller amount will lower the level of sugar in the blood. If a person is not sensitive to insulin it means she or he will need more insulin to lower the level of sugar in the blood. When a person needs more insulin to lower blood sugar, they are more resistant to insulin.

Insulin-to-carb ratio

A formula you use to match the dose of insulin to the amount of carbs you eat and drink.

Intermediate-acting insulin

This is a type of basal insulin. It controls blood sugar for about half the day or overnight. This insulin starts working in about 1 to 2 hours. It works best in your body at 4 to 12 hours and then starts fading. How it works is different for each person.

NPH is the generic name of the drug. Humulin [N], Novolin [N], or ReliOn [N] are brand names.

This insulin looks cloudy. You can mix it with regular or rapid-acting insulin in a syringe. See basal insulin, regular insulin and rapid-acting insulin for more information.

Islets of Langerhans

Small islands of cells scattered throughout the pancreas that make hormones. They have beta-cells, which make insulin and alpha cells which make glucagon. Other cells include delta cells, PP cells and Epsilon cells which make other hormones.

Ketoacidosis—See Diabetic ketoacidosis

Ketones

The body releases these acids when body fat breaks down.

Ketones can build up to dangerous levels in the absence of insulin. This is because the body is not able to break down sugar as fuel.

A urine or a blood test can measure them. A urine dip stick is usually used.

Lancet

A spring-loaded device that you use to prick the skin with a small needle. You do this to get a drop of blood to check your blood sugar.

Lipodystrophy

This is when the fat tissue below the skin becomes swollen, hard or forms dimples. It also limits the body from absorbing insulin if you inject in that area.

Giving yourself many shots into the same area of skin or putting the pump cannula in the same site time after time often causes this.

Lantus

This is a brand drug name of one kind of basal insulin. The generic name for Lantus is glargine. This long-acting basal insulin drug comes in one strength written as U100. See basal insulin and long-acting insulin for more information.

Levemir

This is a brand drug name of one kind of basal insulin. The generic name for Levemir is detemir. This long-acting basal insulin drug comes in one strength written as U100. See basal insulin and long-acting insulin for more information.

Lispro

This is a generic drug name of one kind of rapid-acting insulin. The brand name for lispro is Humalog. See rapid-acting insulin for more information.

Long-acting insulin

This type of basal insulin controls blood sugar consistently for an entire day or longer. After injecting, it begins working many hours and can stay in the bloodstream up to 42 hours. How long it works can be different for different people. It may start weakening a few hours earlier for some, while it may work a few hours longer for others. It comes in different strengths shown as U100, U200 and U300.

See basal insulin to learn more.

Long-acting insulins are:

Generic name	Brand name
Degludec U100	Tresiba U100
Degludec U200	Tresiba U200
Detemir U100	Levemir
Glargine U100	Lantus or Basaglar
Glargine U300	Toujeo

Medical insurance or health insurance

This is a plan that a person signs up for that pays for some or all the costs of medical and surgical care. These plans differ from state to state. Sometimes people must buy their own insurance. Other times they get it from their job or the government. Government plans include Medicare and Medicaid. In some states, the plan may have its own name. For instance, in California it is Medi-Cal.

Multiple daily injections (MDI)

This is a schedule where you give yourself many insulin shots each day. In most cases, you use a long-acting insulin along with shots of rapid-acting insulin before each meal or snack. Some people also use intermediate-acting insulin. See long-acting, intermediate-acting and rapid-acting insulin for more information.

Novolin [N]

This is a brand drug name of one kind of intermediate-acting insulin. The generic name for Novolin [N] is NPH. See intermediate-acting insulin for more information. It is a cloudy insulin.

Novolin [R]

This is a brand drug name of one kind of short-acting insulin. The generic name for Novolin [R] is Insulin Regular. See short-acting insulin for more information.

NPH

This is a generic drug name of an intermediate-acting insulin. The brand names for NPH are Humulin [N], Novolin [N] or ReliOn [N]. See intermediate-acting insulin for more information.

Occlusion

The infusion set or infusion site clogs or blocks. This can stop or slow insulin delivery.

In most cases, an occlusion happens when the cannula gets pinched, kinked or dislodged. The cannula blocks when insulin crystals form.

An occlusion can be partial. That means it only reduces, but does not stop the flow of insulin. Or it can be complete. That means no insulin gets through the tubing.

Pancreas

This gland is near the stomach. It is deep in the center of the body. It releases insulin and other hormones. It also releases digestive enzymes.

Pharmacist

This health care professional prepares and gives medicine to people. She or he also gives information on medicines.

Pre-mixed insulin

In most cases, people with diabetes take these two or three times a day before a meal. They are insulins where a shorter and longer acting insulin mixed. In most cases, they look cloudy. The numbers after the name describe how much long-acting and short-acting insulin is in the mix. They have many names, including:

- Humulin 70/30 (70% long acting/30% short acting insulin)
- Novolin 70/30 (70% long acting/30% short acting insulin)
- Novolog 70/30 (70% long acting/30% rapid acting insulin)
- Humulin 50/50 (50% long acting/50% short acting insulin)
- Humalog mix 75/25 (75% long acting/25% rapid acting insulin)

Proteins

These are one of the three main parts of foods along with carbohydrates and fats. Proteins are made of amino acids. Foods like milk, meat, fish, and eggs have protein.

The body burns proteins more slowly than fats or carbohydrates. There are four calories per gram of protein.

Rapid-acting insulin

If you give yourself shots, you will give both long-acting insulin and short or rapid-acting insulin. The rapid-acting insulin covers insulin needs for meals. You give yourself a shot at the same time you eat.

If you use a pump, you only use rapid acting insulin. The pump gives out rapid-acting insulin in small amounts on an ongoing basis. You also program your pump to give you a bolus of insulin for meals. See long acting insulin and bolus for more information.

Rapid acting insulins are:

Generic name	Brand name
Lispro	Humalog
Aspart	Novolog
Glulisine	Apidra

Reservoir, syringe, cartridge

This container holds the fast-acting insulin inside a pump.

Self-management

In diabetes, this means the ongoing process of managing diabetes. It includes when you:

- Plan meals
- Plan physical activity
- Check blood sugar
- Take diabetes medicines
- Handle diabetes when you are sick
- Handle low and high blood sugar
- Manage your diabetes on trips

People with diabetes design their own self-management treatment plan. They do this with the support of their diabetes team. This includes doctors, nurses, dietitians, pharmacists and others.

Sensitivity factor

This is the amount that a single unit of insulin lowers the blood sugar level in a person. Often this is first set at 50. But based on how a person reacts to insulin it can change.

A lower number, such as 25, means that the person is less sensitive to insulin. A higher number, such as 75, means that the person is more sensitive to insulin.

Sharps container

This is a container where you get rid of used needles and syringes. It is often made of hard plastic so that needles cannot poke through.

Self-monitoring of blood glucose (SMBG)

This is when you check your blood sugar with a blood sugar meter.

Short-acting insulin

Short-acting insulin covers insulin needs for meals. You give yourself a shot about 30 minutes before you eat. Short-acting insulin brand names are Humulin [R], Novolin [R] or ReliOn [R]. The generic name is regular insulin.

Starch

This is a type of complex carbohydrate. Some examples are bread, pasta and rice.

Sugar

A kind of carbohydrate that most often has a sweet taste. This includes glucose, fructose and sucrose. In the diabetes world, the word sugar is often used instead of glucose. Blood glucose and blood sugar mean the same thing.

Sugar alcohol

This is a sugar substitute. It has simple sugars with an alcohol molecule attached to them. This lowers the calorie content. It also delays the effect on blood sugar levels.

Syringe

This is a device used to inject medication or other liquids into body tissues. The syringe for insulin has a hollow plastic tube with a plunger inside. It also has a needle on the end.

Team management

This is an approach to treat diabetes where a team provides medical care. See Diabetes team for more information.

Total daily dose (TDD)

The total amount of insulin a person uses in a day. It means adding all the insulin doses: faster and slower acting insulin together. You use the TDD to help figure out the basal rate, carb factor and correction factor.

Tresiba

This is a brand drug name of one kind of basal insulin. The generic name for Tresiba is degludec. This long-acting basal insulin drug comes in two strengths written as either U100 or U200. See basal insulin and long-acting insulin for more information.

Toujeo

This is a brand drug name of one kind of basal insulin. The generic name for Toujeo is glargine. This long-acting basal insulin drug comes in one strength written as U300. See basal insulin and long-acting insulin for more information.

Type 1 Diabetes

In Type 1 Diabetes, the pancreas makes little or no insulin. This is because the beta cells in the body that make insulin are destroyed.

It is an autoimmune disease. This is caused by a defect where the body's internal defense system attacks a part of the body itself.

Most often, this type of diabetes appears suddenly. It is more common in people younger than 30. But it can appear at any age.

The ways to treat it are:

- Give daily insulin shots or use an insulin pump
- Count carbohydrates
- Exercise regularly
- Self-monitor blood sugar levels each day through finger sticks or by using a continuous glucose monitoring (CGM).

Units of insulin

This is the basic measure of insulin. U-100 insulin means 100 units of insulin per milliliter (mL) or cubic centimeter (cc) of solution.

It is a way to describe the concentration of insulin. In the United States, there are U100, U200, U300 and U500 insulins.